

Gravitational field of a slowly rotating black hole with phantom global monopole

Songbai Chen^{*}, Jiliang Jing[†]

*Institute of Physics and Department of Physics, Hunan Normal University,
Changsha, Hunan 410081, People's Republic of China
Key Laboratory of Low Dimensional Quantum Structures
and Quantum Control of Ministry of Education, Hunan Normal University,
Changsha, Hunan 410081, People's Republic of China*

Abstract

We present a slowly rotating black hole with phantom global monopole by solving Einstein's field equation and find that presence of global monopole changes the structure of black hole. The metric coefficient $g_{t\phi}$ contains hypergeometric function of the polar coordinate r , which is more complex than that in the usual slowly rotating black hole. The energy scale of symmetry breaking η affects the black hole horizon and a deficit solid angle. Especially, the solid angle is surplus rather than deficit for a black hole with the phantom global monopole. We also study the correction originating from the global monopole to the angular velocity of the horizon Ω_H , the Kepler's third law, the innermost stable circular orbit and the radiative efficiency ϵ in the thin accretion disk model. Our results also show that for the phantom black hole the radiative efficiency ϵ is positive only for the case $\eta \leq \eta_c$. The threshold value η_c increases with the rotation parameter a .

PACS numbers: 04.70.Dy, 95.30.Sf, 97.60.Lf

^{*} csb3752@hunnu.edu.cn

[†] jljing@hunnu.edu.cn

I. INTRODUCTION

Phantom field is a special kind of dark energy model with the negative kinetic energy [1], which is applied extensively in cosmology to explain the accelerating expansion of the current Universe [2–7]. Comparing with other dark energy models, the phantom field is more interesting because that the presence of negative kinetic energy results in that the equation of state of the phantom field is less than -1 and then the null energy condition is violated. Although the phantom field owns such exotic properties, it is still not excluded by recent precise observational data [8], which encourages many people to focus on investigating phantom field from various aspects of physics.

Phantom field also exhibits some peculiar properties in the black hole physics. E. Babichev [9] found that the mass of a black hole decreases when it absorbs the phantom dark energy. This means that the cosmic censorship conjecture is challenged severely by a fact that the charge of a Reissner-Nordström-like black hole absorbing the phantom energy will be larger than its mass. We studied the wave dynamics of the phantom scalar perturbation in the Schwarzschild black hole spacetime and that in the late-time evolution the phantom scalar perturbation grows with an exponential rate rather than decays as the usual scalar perturbations [10, 11]. Moreover, we also find that the phantom scalar emission will enhance the Hawking radiation of a black hole [12]. Furthermore, some black hole solutions describing gravity coupled to phantom scalar fields or phantom Maxwell fields have also been found in [13–20]. The thermodynamics and the possibility of phase transitions in these phantom black holes are studied in [16, 17]. The gravitational collapse of a charged scalar field [18] and the light paths [19] are investigated in such kind of spacetimes. Moreover, S. Bolokhov *et al* also study the regular electrically and magnetically charged black hole with a phantom scalar in [20]. These investigations could help us to get a deeper understand about dark energy and black hole physics.

A global monopole is one of the topological defects which could be formed during phase transitions in the evolution of the early Universe. The metric describing a static black hole with a global monopole was obtained by Barriola and Vilenkin [21], which arises from the breaking of global $SO(3)$ symmetry of a triplet scalar field in a Schwarzschild background. Due to the presence of the global monopole, the black hole owns different topological structure from that of the Schwarzschild black hole. The physical properties of the black hole with a global monopole have been studied extensively in recent years [22–25].

The main purpose of this paper is to study the gravitational field of phantom global monopole arising from a triplet scalar field with negative kinetic energy and to see how the energy scale of symmetry breaking

η influences the structure of black hole, the Kepler's third law, the innermost stable circular orbit and the radiative efficiency ϵ in the thin accretion disk model. Moreover, we will explore how it differs from that in the black hole with ordinary global monopole.

The paper is organized as follows: in the following section we will construct a static and spherical symmetric solution of a phantom global monopole from a triplet scalar field with negative kinetic energy, and then study the effect of the parameter η on the black hole. In Sec.III, we obtain a slowly rotating black hole with phantom global monopole by solving Einstein's field equation and find that presence of global monopole make the metric coefficient of black hole more complex. In Sec.IV, we will focus on investigating the effects of the parameter η on the Kepler's third law, the innermost stable circular orbit and the radiative efficiency ϵ in the thin accretion disk model. We end the paper with a summary.

II. A STATIC AND SPHERICAL SYMMETRIC BLACK HOLE WITH PHANTOM GLOBAL MONOPOLE

Let us now first study a static and spherical symmetric black hole with phantom global monopole formed by spontaneous symmetry breaking of a triplet of phantom scalar fields with a global symmetry group $O(3)$. The action giving rise to the phantom global monopole is

$$S = \int \sqrt{-g} d^4x [R - \frac{\xi}{2} \partial^\mu \psi^a \partial_\mu \psi^i - \frac{\lambda}{4} (\psi^i \psi^i - \eta^2)^2], \quad (1)$$

where ψ^i is a triplet of scalar field with $i = 1, 2, 3$, η is the energy scale of symmetry breaking and λ is a constant. The coupling constant ξ in the kinetic term takes the value $\xi = 1$ corresponds to the case of the ordinary global monopole originating from the scalar field with the positive kinetic energy [21]. As the coupling constant $\xi = -1$, the kinetic energy of the scalar field is negative and then the phantom global monopole is formed.

Following in Ref.[21], we can take ansatz describing a monopole as

$$\psi^a = \frac{\eta f(r) x^i}{r}, \quad (2)$$

where $x^i x^i = r^2$. Equipping with the general static and spherical symmetric metric

$$ds^2 = -B(r)dt^2 + A(r)dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2, \quad (3)$$

one can find that the field equations for ψ^i can be reduced to a single equation for $f(r)$

$$\frac{\xi f''}{A} + \left[\frac{2}{Ar} + \frac{1}{2B} \left(\frac{B}{A} \right)' \right] \xi f' - \frac{2\xi f}{r^2} - \lambda \eta^2 f(f^2 - 1) = 0. \quad (4)$$

Moreover, the energy-momentum tensor for the spacetime with a global monopole can be expressed as

$$T_t^t = \left[\frac{\eta^2 f'^2}{2A} + \frac{\eta^2 f^2}{r^2} \right] \xi + \frac{\lambda}{4} \eta^4 (f^2 - 1)^2, \quad (5)$$

$$T_r^r = \left[-\frac{\eta^2 f'^2}{2A} + \frac{\eta^2 f^2}{r^2} \right] \xi + \frac{\lambda}{4} \eta^4 (f^2 - 1)^2, \quad (6)$$

$$T_\theta^\theta = T_\phi^\phi = \frac{\xi \eta^2 f'^2}{2A} + \frac{\lambda}{4} \eta^4 (f^2 - 1)^2. \quad (7)$$

Similarly, as in Ref.[21], one can take an approximation $f(r) = 1$ outside the core due to a fact that $f(r)$ grows linearly when $r < (\eta\sqrt{\lambda})^{-1}$ and tends exponentially to unity as soon as $r > (\eta\sqrt{\lambda})^{-1}$. With this approximation, we can obtain a solution of the Einstein equations

$$B = A^{-1} = 1 - 8\pi\xi\eta^2 - \frac{2M}{r}, \quad (8)$$

where M is an integrate constant. Obviously, the radius of event horizon is $r_H = 2M/(1 - 8\pi\xi\eta^2)$. Here we must point that it is possible that the radius of event horizon r_H is larger than the monopole's core $\delta \sim (\eta\sqrt{\lambda})^{-1}$ if the coupling constant $\lambda \gg \frac{(1-8\pi\xi\eta^2)^2}{4M^2\eta^2}$. In this case, the metric (3) can describe the geometry near the horizon in the spacetime with global monopole. With the increase of the energy scale of symmetry breaking η , one can find that the radius of event horizon r_H increases for a Schwarzschild black hole with the ordinary global monopole (S_+), but decreases for a Schwarzschild black hole with the phantom global monopole (S_-). Thus, comparing with the system S_+ , one can find that the system S_- possesses the higher Hawking temperature and the lower entropy. In the low energy limit, the luminosity of Hawking radiation of a spherical symmetric black hole can be approximated as $L = \frac{2\pi^3 r_H^2}{15} T_H^4 \propto (1 - 8\pi\xi\eta^2)^4 / r_H^2$, which tells us that the energy scale of symmetry breaking η enhances Hawking radiation for the system S_- , but it decreases Hawking radiation in the system S_+ . The presence of the phantom field enhances the Hawking emission of Kerr black hole are also found in [12].

Introducing the following transformations

$$t \rightarrow (1 - 8\pi\xi\eta^2)^{1/2} t, \quad r \rightarrow (1 - 8\pi\xi\eta^2)^{1/2} r, \quad M \rightarrow (1 - 8\pi\xi\eta^2)^{3/2} M, \quad (9)$$

one can rewrite the metric (3) with the functions (8) as

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + (1 - 8\pi\xi\eta^2) r^2 (d\theta^2 + \sin^2 \theta d\phi^2). \quad (10)$$

It is clear that there exists a deficit solid angle $(1 - 8\pi\eta^2)$ for the system S_+ . However, for the system S_- (i.e., $\xi = -1$), one can find that the solid angle becomes $(1 + 8\pi\eta^2)$, which is surplus rather than deficit. This implies that the topological properties of a spacetime with the phantom global monopole is different from that of with a ordinary global monopole.

III. A SLOWLY ROTATING BLACK HOLE WITH PHANTOM GLOBAL MONOPOLE

In this section, we first obtain the metric for a slowly rotating black hole with phantom global monopole by solving Einstein's field equation. And then, we will study the properties of the black hole spacetime.

From the static and spherical symmetric solution (3) with the metric function (8), we can assume the metric has a form for a slowly rotating black hole with phantom global monopole

$$ds^2 = -U(r)dt^2 + \frac{1}{U(r)}dr^2 - 2F(r, \theta)adtd\phi + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (11)$$

where a is a parameter associated with its angular momentum. Moreover, we assume that in the slowly rotating spacetime (11) the ansatz describing a monopole is

$$\psi^a = \frac{\eta f(r)x^i}{r}. \quad (12)$$

It is easy to find that the field equations for ψ^i can also be reduced to a single equation for $f(r)$

$$\xi f''U(r) + \left[\frac{2U(r)}{r} + U(r)' \right] \xi f' - \frac{2\xi f}{r^2} - \lambda \eta^2 f(f^2 - 1) = 0, \quad (13)$$

which is similar to that in the static and spherical symmetric spacetime (3). It is not surprising since the triplet scalar field ψ^i does not depend on the time coordinate t .

Inserting the metric (11) and the triplet scalar (12) into the Einstein field equation, we find that the non-vanishing components of field equation can be expanded to first order in the angular momentum parameter a as

$$tt: \quad \frac{U(r)}{r^2} \left[U'(r)r + U(r) - 1 \right] + 8\pi U(r) \left[\left(\frac{\eta^2 f'^2 U(r)}{2} + \frac{\eta^2 f^2}{r^2} \right) \xi + \frac{\lambda}{4} \eta^4 (f^2 - 1)^2 \right] = 0 + \mathcal{O}(a^2), \quad (14)$$

$$rr: \quad \frac{1}{U(r)r^2} \left[U'(r)r + U(r) - 1 \right] + \frac{8\pi}{U(r)} \left[\left(-\frac{\eta^2 f'^2 U(r)}{2} + \frac{\eta^2 f^2}{r^2} \right) \xi + \frac{\lambda}{4} \eta^4 (f^2 - 1)^2 \right] = 0 + \mathcal{O}(a^2), \quad (15)$$

$$\theta\theta: \quad -\frac{r}{2} \left[U''(r)r + 2U'(r) \right] + 8\pi r^2 \left[\frac{\xi \eta^2 f'^2 U(r)}{2} + \frac{\lambda}{4} \eta^4 (f^2 - 1)^2 \right] = 0 + \mathcal{O}(a^2), \quad (16)$$

$$\phi\phi: \quad -\frac{r}{2} \sin^2 \theta \left[U''(r)r + 2U'(r) \right] + 8\pi r^2 \left[\frac{\xi \eta^2 f'^2 U(r)}{2} + \frac{\lambda}{4} \eta^4 (f^2 - 1)^2 \right] \sin^2 \theta = 0 + \mathcal{O}(a^2), \quad (17)$$

$$t\phi: \quad \frac{1}{2r^2} \left\{ r^2 U(r) \frac{\partial^2 F(r, \theta)}{\partial r^2} - F(r, \theta) \left[r^2 U''(r) + 2rU'(r) + 2U(r) \right] + 2F(r, \theta) + \frac{\partial^2 F(r, \theta)}{\partial \theta^2} - \frac{\partial F(r, \theta)}{\partial \theta} \cot \theta \right\} \\ - 8\pi F(r, \theta) \left[\left(\frac{\eta^2 f'^2 U(r)}{2} + \frac{\eta^2 f^2}{r^2} \right) \xi + \frac{\lambda}{4} \eta^4 (f^2 - 1)^2 \right] = 0 + \mathcal{O}(a^2). \quad (18)$$

Solving the Einstein equations (14)-(17) with the approximation $f(r) = 1$ outside the core, we can obtain the metric coefficient

$$U(r) = 1 - 8\pi \xi \eta^2 - \frac{2M}{r}. \quad (19)$$

Separating $F(r, \theta) = h(r)\Theta(\theta)$, we can obtain the equation for the angular part

$$\frac{d^2\Theta(\theta)}{d\theta^2} - \frac{d\Theta(\theta)}{d\theta} \cot \theta = \lambda\Theta(\theta). \quad (20)$$

In order to that the coefficient $g_{t\phi}$ can be reduced to that in the slowly rotating black hole without the global monopole, here we set $\lambda = -2$ and find that $\Theta(\theta) = \sin^2(\theta)$ in this case. And then the radial part of Eq.(18) becomes

$$r^2 U(r) \frac{d^2 h(r)}{dr^2} - 2h(r)U(r) - 16\pi\xi\eta^2 h(r) = 0. \quad (21)$$

Substituting Eq. (19) into the above radial equation, we obtain (see in appendix)

$$h(r) = \left[\frac{2M}{(1-b)r} \right]^{\frac{1}{2}(\sqrt{\frac{9-b}{1-b}}-1)} {}_2F_1 \left[\frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} - 3 \right), \frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} + 3 \right), \sqrt{\frac{9-b}{1-b}} + 1, \frac{2M}{(1-b)r} \right], \quad (22)$$

where $b = 8\pi\xi\eta^2$ and ${}_2F_1[a_1, b_1, c_1; x]$ is the hypergeometric function. As a usual slowly rotating black hole, the horizon of black hole (11) is given by the zeros of the function $U(r) = (g_{rr})^{-1}$, i.e., $r_H = \frac{2M}{1-b}$, which is the same as that in the static and spherical symmetric case (3). The mainly reason is that we here expand the metric only to first order in the angular momentum parameter a .

From Eq. (22), it is obvious to see that due to the presence of the global monopole the form of the metric coefficient $g_{t\phi}$ becomes more complicated in the slowly rotating black hole (11). The dependence of $h(r)$ on the parameter η is shown in Fig. (1), which tells us that for the slowly rotating black hole with the ordinary global monopole (SR_+) the function $h(r)$ increases with the parameter η near the horizon, but decreases at the far field region with the larger value of r . For a slowly rotating black hole with the phantom global monopole (SR_-), the behavior of $h(r)$ is just the opposite. In a word, the dependence of the function $h(r)$ on the η near the horizon is different that in the far field region in these two global monopole cases. As the parameter $b \rightarrow 0$, one can find that $h(r) \rightarrow \frac{2M}{r}$, which recovers that of a slowly rotating black hole without global monopole. When the rotation parameter a vanishes, one can get the previous solution of a static and spherical symmetric black hole with phantom global monopole (3).

For a rotating black hole, one of the important quantities is the angular velocity of the horizon Ω_H , which affects the region where the super-radiance occurs in the black hole background. In the spacetime of a slowly rotating black hole with global monopole, the angular velocity of the horizon Ω_H can be expressed as

$$\Omega_H = - \frac{g_{t\phi}}{g_{\phi\phi}} \Big|_{r=r_H} = \frac{a(1-b)^2}{r_H^2} \frac{\Gamma[\sqrt{\frac{9-b}{1-b}} + 1]}{4\Gamma[\frac{1}{2}(\sqrt{\frac{9-b}{1-b}} - 1)]\Gamma[\frac{1}{2}(\sqrt{\frac{9-b}{1-b}} + 5)]}, \quad (23)$$




FIG. 1: The dependence of $h(r)$ on the parameter η . The left and right panels are for the systems SR_+ and SR_- , respectively. Here we set $M = 1$.

which depends on the energy scale of symmetry breaking η . We plot the change of the angular velocity Ω_H with the parameter η in Fig.(2). For the system SR_+ , we find that the angular velocity Ω_H first decreases slowly and then increases rapidly with the increase of η . There exists a minimum for Ω_H at where $\eta = 0.1725$ (i.e., $b = 0.7477$). This means that there exist a minimum region for the occurrence of the super-radiance in the black hole with the ordinary global monopole for fixed a . For the the system SR_- , Ω_H increases monotonically with the energy scale of symmetry breaking η .




FIG. 2: The change of the angular velocity Ω_H with the parameter η . The left and right panels are for the systems SR_+ and SR_- , respectively. Here we set $M = 1$ and $a = 0.2$.

Making the same transformations (9) as in the Sec.II, one find that the area of the horizon becomes $A_H = br_H^2$. It means that for the system SR_- , the solid angle is surplus rather than deficit, which is similar to that in the static and spherical symmetric case (3).

IV. KEPLER'S THIRD LAW AND THE INNERMOST STABLE CIRCULAR ORBIT

In this section, we will focus on how the symmetry breaking scale η of global monopole affects the Kepler's third law and the innermost stable circular orbit (ISCO) in this slowly rotating black hole.

In the stationary and axially symmetric spacetime, one can find that the timelike geodesics take the form

$$\dot{t} = \frac{Eg_{\phi\phi} - L_z g_{t\phi}}{g_{t\phi}^2 + g_{tt}g_{\phi\phi}}, \quad (24)$$

$$\dot{\phi} = \frac{Eg_{t\phi} + L_z g_{tt}}{g_{t\phi}^2 + g_{tt}g_{\phi\phi}}, \quad (25)$$

$$g_{rr}\dot{r}^2 + g_{\theta\theta}\dot{\theta}^2 = V_{eff}(r, \theta; E, L_z), \quad (26)$$

with the effective potential

$$V_{eff}(r) = \frac{E^2 g_{\phi\phi} - 2EL_z g_{t\phi} - L_z^2 g_{tt}}{g_{t\phi}^2 + g_{tt}g_{\phi\phi}} - 1, \quad (27)$$

where the overhead dot stands for a derivative with respect to the affine parameter. The constants E and L_z correspond to the conserved energy and the (z -component of) orbital angular momentum of the particle, respectively.

For simplicity, we set the orbits on the equatorial plane. With the restriction that $\theta = \pi/2$, one can find that for the stable circular orbit in the equatorial plane, the effective potential $V_{eff}(r)$ must obey

$$V_{eff}(r) = 0, \quad \frac{dV_{eff}(r)}{dr} = 0. \quad (28)$$

Solving above equations, one can obtain

$$\begin{aligned} E &= \frac{g_{tt} + g_{t\phi}\Omega}{\sqrt{g_{tt} + 2g_{t\phi}\Omega - g_{\phi\phi}\Omega^2}}, \\ L_z &= \frac{-g_{t\phi} + g_{\phi\phi}\Omega}{\sqrt{g_{tt} + 2g_{t\phi}\Omega - g_{\phi\phi}\Omega^2}}, \\ \Omega &= \frac{d\phi}{dt} = \frac{g_{t\phi,r} + \sqrt{(g_{t\phi,r})^2 + g_{tt,r}g_{\phi\phi,r}}}{g_{\phi\phi,r}}, \end{aligned} \quad (29)$$

where Ω is the angular velocity of particle moving in the orbits. From Eq.(29), one can obtain Kepler's third law in the slowly-rotating black-hole spacetime with the global monopole

$$\begin{aligned} T^2 &= \frac{4\pi^2 R^3}{M} \left[1 + \frac{4a}{(1-b)^2(\sqrt{\frac{9-b}{1-b}} + 1)M^{1/2}R^{3/2}} \left[\frac{2M}{(1-b)R} \right]^{\frac{1}{2}(\sqrt{\frac{9-b}{1-b}} - 1)} \left\{ (1-b)R \times \right. \right. \\ &\quad \left. \left. {}_2F_1 \left[\frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} - 3 \right), \frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} + 3 \right), \sqrt{\frac{9-b}{1-b}} + 1, \frac{2M}{(1-b)R} \right] \right. \right. \\ &\quad \left. \left. + bM {}_2F_1 \left[\frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} - 1 \right), \frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} + 5 \right), \sqrt{\frac{9-b}{1-b}} + 2, \frac{2M}{(1-b)R} \right] \right\} + \mathcal{O}(a^2) \right], \end{aligned} \quad (30)$$

where T is the orbital period and R is the radius of the circular orbit. The later terms in the right hand side is the correction by the a and the symmetry breaking scale η of global monopole. From Eq.(30), we find that for fixed R the η affects the orbital period T only in the case with nonzero rotation parameter a . In Fig. (3), we present the change of the corrected term $\Delta T^2 = \frac{4\pi^2 T^2 R^3}{M} - 1$ with the parameter η in this spacetime. It is shown that the absolute value $|\Delta T^2|$ increases with the scale η for the system SR_+ , but decreases with η in the system SR_- . Moreover, we also find that the presence of the global monopole make the orbital period T increase for a prograde particle (i.e., $a > 0$) and decrease for a retrograde one (i.e., $a < 0$).




FIG. 3: The change of the corrected term $\Delta T^2 = \frac{4\pi^2 T^2 R^3}{M} - 1$ in the orbital period T with the parameter η . The solid and dashed lines are for the systems SR_+ and SR_- , respectively. Here we set $M = 1$, $a = 0.2$ and $R = 10M$.

The innermost stable circular orbit (ISCO) of the particle around the black hole is given by the condition $V_{eff,rr} = 0$. For a slow rotating black hole with the global monopole (11), we obtain

$$\begin{aligned}
 r_{ISCO} = & \frac{6M}{1-b} - \frac{4a\sqrt{2} 3^{-2-\frac{1}{2}} \sqrt{\frac{9-b}{1-b}}}{(1-b)^{5/2} (\sqrt{\frac{9-b}{1-b}} + 1) (\sqrt{\frac{9-b}{1-b}} + 2)} \left\{ 9 \left[\sqrt{(9-b)(10-3b)} + (3b^2 - 25b + 30) \right] \times \right. \\
 & {}_2F_1 \left[\frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} - 3 \right), \frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} + 3 \right), \sqrt{\frac{9-b}{1-b}} + 1, \frac{1}{3} \right] + 3b \left[5\sqrt{(9-b)(1-b)} + (15-7b) \right] \times \\
 & {}_2F_1 \left[\frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} - 1 \right), \frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} + 5 \right), \sqrt{\frac{9-b}{1-b}} + 2, \frac{1}{3} \right] + b \left[\sqrt{(9-b)(1-b)} + (1+b) \right] \times \\
 & \left. {}_2F_1 \left[\frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} + 1 \right), \frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} + 7 \right), \sqrt{\frac{9-b}{1-b}} + 3, \frac{1}{3} \right] \right\} + \mathcal{O}(a^2), \quad (31)
 \end{aligned}$$

Obviously, the ISCO radius r_{ISCO} decreases with the rotation parameter a , which is similar to that in the usual Kerr black hole. In Fig.(4), we set $M = 1$ and plotted the variety of the ISCO radius r_{ISCO} with the parameter η in the slowly rotating black hole spacetime with the global monopole. It is shown that the ISCO radius r_{ISCO} increases with the scale η for the system SR_+ , but decreases with η in the system SR_- . Moreover, we find that the ISCO radius in the system SR_+ is larger than that in the system SR_- for fixed a .




FIG. 4: The change of the ISCO radius r_{ISCO} with the parameters η and a . The dashed and solid lines are for the systems SR_+ and SR_- , respectively. The thin and thick lines correspond to the cases with $a = -0.2$ and $a = 0.2$. Here we set $M = 1$.

Let us now compute the effect of the symmetry breaking scale η on the radiative efficiency ϵ in the thin accretion disk model, which is defined by

$$\epsilon = 1 - E(r_{ISCO}). \quad (32)$$

This quantity corresponds to the maximum fraction of energy being radiated when a test particle accretes into a central black hole. For Schwarzschild and extremal Kerr black holes, $\epsilon \sim 0.06$ and $\epsilon \sim 0.42$, respectively.

For a slowly rotating black hole with the global monopole, the radiative efficiency ϵ can be expressed as

$$\begin{aligned} \epsilon = & 1 - \frac{2\sqrt{2}}{3}\sqrt{1-b} + \frac{a}{M(\sqrt{\frac{9-b}{1-b}}+1)} 3^{-2-\frac{1}{2}\sqrt{\frac{9-b}{1-b}}} \left\{ \sqrt{(9-b)} \left[\sqrt{(9-b)} + \sqrt{(1-b)} \right] \times \right. \\ & {}_2F_1 \left[\frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} - 3 \right), \frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} + 3 \right), \sqrt{\frac{9-b}{1-b}} + 1, \frac{1}{3} \right] + \frac{2b}{3} \times \\ & \left. {}_2F_1 \left[\frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} - 1 \right), \frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} + 5 \right), \sqrt{\frac{9-b}{1-b}} + 2, \frac{1}{3} \right] \right\} + \mathcal{O}(a^2), \end{aligned} \quad (33)$$

It is clear that the radiative efficiency ϵ increases with the rotation parameter a in these two global monopole black holes. The effect of the symmetry breaking scale η on the radiative efficiency ϵ is shown in Fig. (5). It tells us that with the scale η the radiative efficiency ϵ increases for the system SR_+ , but decreases for the system SR_- . Moreover, we also note that for the phantom black hole the radiative efficiency ϵ is positive only for the case $\eta \leq \eta_c$. This could be explained by a fact that for a phantom black hole with $\eta \geq \eta_c$, its capability of capturing particle could become so weak that the accreted matter around it is very dilute, which could lead to that the radiation can not be generated because of lacking of the enough stress and dynamic




FIG. 5: The change of the radiative efficiency ϵ with the parameter η and a . The dashed and solid lines are for the systems SR_+ and SR_- , respectively. The thin and thick lines correspond to the cases with $a = -0.2$ and $a = 0.2$. Here we set $M = 1$.

friction in the accretion disk model. The critical value η_c can be approximated as

$$\eta_c = 0.0705 + 0.0227 \frac{a}{M} + \mathcal{O}(a^2). \quad (34)$$

It means that the critical value η_c increases with the rotation parameter a , which is also shown in Fig.(6).




FIG. 6: The change of the critical value η_c with the rotation parameter a for the system SR_- . Here we set $M = 1$.

V. SUMMARY

In this paper we present firstly a four-dimensional spherical symmetric black hole with phantom global monopole and find that the scale of symmetry breaking η affects the radius of the black hole horizon and a deficit solid angle. For the system SR_- , the solid angle is surplus rather than deficit as in the systems SR_+ . Then, we obtain a slowly rotating black hole solution with global monopole by solving Einstein's field equation.

We find that presence of global monopole makes the metric coefficient $g_{t\phi}$ contain the hypergeometric function of the polar coordinate r , which is more complex than that in the usual slowly rotating black hole. We study the property of the angular velocity of the horizon Ω_H , which is connected with the region where the super-radiance occurs in the black hole background. With the increase of η , the angular velocity Ω_H first decreases slowly and then increases rapidly for the system SR_+ , but increases monotonically with η for the system SR_- .

We also analyze the effects of the scale of symmetry breaking η on the Kepler's third law, the innermost stable circular orbit and the radiative efficiency ϵ in the thin accretion disk model. Our results show that only in the case with nonzero rotation parameter a the orbital period T depends on the scale of symmetry breaking η for fixed orbital radius R . The presence of the global monopole make the orbital period T increase for a prograde particle (i.e., $a > 0$) and decrease for a retrograde one (i.e., $a < 0$). All of the absolute value of the corrected term to Kepler's Third Law (i.e., $|\Delta T^2|$), the ISCO radius r_{ISCO} and the radiative efficiency ϵ in the thin accretion disk model increase with the scale η for the system SR_+ , but decrease with η in the system SR_- . Moreover, we also find that for the phantom black hole the radiative efficiency ϵ is positive only for the case $\eta \leq \eta_c$. The threshold value η_c increases with the rotation parameter a .

VI. ACKNOWLEDGMENTS

This work was partially supported by the National Natural Science Foundation of China under Grant No.11275065, the NCET under Grant No.10-0165, the PCSIRT under Grant No. IRT0964, the Hunan Provincial Natural Science Foundation of China (11JJ7001) and the construct program of key disciplines in Hunan Province. J. Jing's work was partially supported by the National Natural Science Foundation of China under Grant Nos. 11175065, 10935013; 973 Program Grant No. 2010CB833004.

Appendix A: The form of $h(r)$

Here we present the form of $h(r)$ by solving the radial equation (21). Defining $z = \frac{2M}{(1-b)r}$, the radial equation (21) can be expressed as

$$z(1-z)\frac{d^2h(z)}{dz^2} + 2(1-z)\frac{dh(z)}{dz} + 2\left[1 - \frac{b}{(1-b)z}\right]h(z) = 0. \quad (\text{A1})$$

Employing the transformation $h(z) = z^\alpha(1-z)^\beta H(z)$, we can write Eq. (A1) into the standard form of the hypergeometric equation

$$z(1-z)\frac{d^2H(z)}{dz^2} + [c - (1+a_1+b_1)z]\frac{dH(z)}{dz} - a_1b_1H(z) = 0, \quad (\text{A2})$$

with

$$c_1 = 2 + 2\alpha, \quad a_1 = \alpha + \beta + 2, \quad b_1 = \alpha + \beta - 1. \quad (\text{A3})$$

Because of the constraint from coefficient of $H(z)$, the power coefficients α and β must satisfy the second-order algebraic equations

$$\beta = 0, \quad \alpha(\alpha + 1) - \frac{2}{1-b} = 0. \quad (\text{A4})$$

Considering the asymptotical behavior $h(r)$ at spatial infinite $r \rightarrow \infty$, we choose $\alpha = \frac{1}{2}[\sqrt{\frac{9-b}{1-b}} - 1]$. Then, the function $h(r)$ has the form

$$h(r) = \left[\frac{2M}{(1-b)r} \right]^{\frac{1}{2}(\sqrt{\frac{9-b}{1-b}}-1)} {}_2F_1 \left[\frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} - 3 \right), \frac{1}{2} \left(\sqrt{\frac{9-b}{1-b}} + 3 \right), \sqrt{\frac{9-b}{1-b}} + 1, \frac{2M}{(1-b)r} \right]. \quad (\text{A5})$$

-
- [1] Caldwell R R 2002 *Phys. Lett. B* **545**, 23 (arXiv: astro-ph/9908168)
 - [2] McInnes B 2002 *J. High Energy Phys.* **08** 029 (arXiv: hep-th/0112066)
 - [3] Nojiri S and Odintsov S D 2003 *Phys. Lett. B* **562** 147(arXiv:hep-th/0303117)
 - [4] Chimento L P and Lazkoz R 2003 *Phys. Rev. Lett.* **91** 211301(arXiv:gr-qc/0307111)
 - [5] Boisseau B, Esposito-Farese G, Polarski D and Starobinsky A A 2000 *Phys. Rev. Lett.* **85** 2236 (arXiv:gr-qc/0001066)
 - [6] Gannouji R, Polarski D, Ranquet A and Starobinsky A A 2006 *J. Cosmol. Astropart. Phys.* **09** 016
 - [7] Caldwell R R, Kamionkowski M and Weinberg N N 2003 *Phys. Rev. Lett.* **91** 071301
Nesseris S and Perivolaropoulos L 2004 *Phys. Rev. D* **70** 123529 (arXiv:astro-ph/0410309)
Nojiri S and Odintsov S D 2003 *Phys. Lett. B* **571** 1 (arXiv:hep-th/0306212)
Singh P, Sami M and Dadhich N 2003 *Phys. Rev. D* **68** 023522 (arXiv: hep-th/0305110)
Hao J G and Li X Z 2004 *Phys. Rev. D* **70** 043529 (arXiv: astro-ph/0309746)
Saridakis E N, Gonzalez-Diaz P F and Siguenza C L 2009 *Class. Quant. Grav.* **26** 165003
 - [8] Melchiorri A, Mersini-Houghton L, Odman C J and Trodden M 2003 *Phys. Rev. D* **68** 043509 (arXiv:astro-ph/0211522)
Ainou M A 2013 *Phys. Rev. D* **87** 024012 (arXiv:1209.5232)
 - [9] Babichev E, Dokuchaev V and Eroshenko Y 2004 *Phys. Rev. Lett.* **93** 021102
 - [10] Chen S, Jing J and Pan Q 2009 *Phys. Lett. B* **670** 276
Chen S, Jing J 2009 *J. High Energy Phys.* **03** 081
 - [11] He X, Wang B, Wu S and Lin C 2009 *Phys. Lett. B* **673** 156
 - [12] Chen S and Jing J 2005 *Class. Quant. Grav.* **22** 4651
 - [13] Gibbons G W and Rasheed D A 1996 *Nucl. Phys. B* **476** 515 (arXiv:hep-th/9604177)
Cl'ement G, Fabris J C and Rodrigues M E 2009 *Phys. Rev. D* **79** 064021 (arXiv:0901.4543)
Azreg-Ainou M, Cl'ement G, Fabris J C and Rodrigues M E 2011 *Phys. Rev. D* **83** 124001 (arXiv:1102.4093).

- [14] Gao C J and Zhang S N 2006 arXiv:hep-th/0604114
- [15] Bronnikov K A and Fabris J C 2006 *Phys. Rev. Lett.* **96** 251101 (arXiv:gr-qc/0511109).
- [16] Rodrigues M E and Oporto Z A A 2012 *Phys. Rev. D* **85** 104022(arXiv:1201.5337)
- [17] Jardim D F,Rodrigues M E and Houndjo M J S 2012 arXiv:1202.2830
- [18] Nakonieczna A, Rogatko M and Moderski R 2012 *Phys. Rev. D* **86** 044043 (arXiv:1209.1203)
- [19] Azreg-Ainou M 2012 arXiv:1209.5232
- [20] Bolokhov S V, Bronnikov K A and Skvortsova M V 2012 *Class. Quant. Grav.* **29**, 245006
- [21] Barriola M and Vilenkin A 1989 *Phys. Rev. Lett.* **63** 341
- [22] Yu H 2002 *Phys. Rev. D* **65** 087502
- [23] Paulo J, Pitelli M and Letelier P 2009 *Phys. Rev. D* **80** 104035
- [24] Chen S and Jing J *Mod. Phys. Lett. A* **23** 359
- [25] Rahaman F, Ghosh P, Kalam M and Gayen K 2005 *Mod. Phys. Lett.A* **20** 1627.